



The birth of a star-forming clump in a disk galaxy at $z \sim 2$

Anita Zanella


with E. Daddi, E. Le Floc'h, F. Bournaud et al.

Spineto, 9th June, 2015

Introduction: observations


Galaxies at $z \sim 2$:

- are gas dominated (Daddi+10, Tacconi+ 10)
- host giant star forming regions = clumps (e.g., Elmegreen+05, 09, Förster-Schreiber+ 06)



Local galaxy
 $d = 15.40$ Mpc

Gas fraction $\sim 5\text{--}10\%$
(Elmegreen+ 13)



High- z galaxy
 $z = 2.05$



Gas fraction $\sim 50\%$
(Wuyts+ 12)

Introduction: observations

Clumps in $z \sim 2$ galaxies:

- mostly identified in *HST* rest-frame UV imaging
- have total masses $\sim 10^{8-9} M_{\odot}$
- size ~ 1 kpc
- have SFR $\sim 20 - 50\%$ of the total SFR of the galaxy (e.g., Genzel+08, Förster-Schreiber+11, Newman+12)

clumps



clumps

Förster-Schreiber+ 11


Introduction: simulations

- At high z : large scale gas inflows feed galaxies with gas (Keres+ 09, Dekel+ 09)
- Due to high gas fraction, violent disk instability fragments disks into giant clumps (Bournaud+ 09)


But which is the fate of giant clumps?

- Do they migrate inward and form the **galaxy bulge**? (Dekel+ 11, Bournaud+ 14)
- Are they disrupted by stellar **feedback** in short **timescales**? (Genel+ 12, Murray+ 10) ?

180Myr



420Myr



Bournaud+ 14

Open questions we would like to answer...

How do clumps form?

Clumps lifetime?

Do clumps form the bulge?


Role of stellar feedback?

Clumps SFE?


...key ingredients we need

spatially resolved probe of **stellar mass distribution**
→ imaging

spatially resolved probe of **star formation distribution**
→ UV, spectroscopy (unique for young ages)



H_{160}



$H\alpha$

Sample

Pointed at CL J1449+0856 cluster (Gobat+ 13)


[OIII] emitting galaxies at $1 \leq z \leq 2$

Observations: WFC3 on board HST

Slitless spectroscopy: G_{141} ($\lambda = 1.1 - 1.7 \mu\text{m}$)

Imaging: near-IR (F140W, F105W)
UVIS (F606W)


→ Spatially resolved [OIII], $H\beta$, [OII]
emission line maps



Slitless spectroscopy: 6.4 arcmin^2

Emission line maps

The case of ID568: **off-nuclear** [OIII], H β and [OII] emissions



GALFIT decomposition: diffuse **disk** + off-nuclear **clump**


Offset **significance** $\sim 8\sigma$

A star forming clump

Discarding the **AGN** hypothesis:

X RAYS: no XMM and Chandra detection

BPT (Baldwin+81): in the SF region (MOIRCS follow up)




Clump equivalent width (**EW**):

$$EW = \frac{F_{\text{line}}}{F_{\text{continuum}}}$$

Upper limits on the continuum flux: simulations

$EW_{[\text{OIII}]} \geq 1700 \text{ \AA} \gg$ typical $EW_{[\text{OIII}]}$ of AGNs



An extremely young SF clump


$Z \sim 0.4 Z_{\odot}$

$Re \leq 0.5$ kpc (unresolved)

Age < 10 Myr

First time robust **age** estimate comparable to the typical **free fall time** in a gas-rich turbulent disk

Starburst99 models



Simulations (led by F. Bournaud)



sSFR clump A = 10x sSFR
other clumps

$t = 0$ birthtime clump A

$t = 12$ Myr observed time
for the M_{\star} and SFR map

other clumps are older
(100 – 300 Myr)

**Initial burst of SF confirmed
by observations**



Do massive clumps exist?

Masses of observed giant clumps are overestimated due to blending caused by insufficient resolution?

For our young clump:







Stellar mass: $M_{\star} \leq 3 \times 10^8 M_{\odot}$

from M/L ratio and simulations

Gas mass estimate: $M_{\text{gas}} \leq 2.5 \times 10^9 M_{\odot}$

from Jeans mass
and simulations

$$M_J \sim \frac{\sigma^4 R_d^2}{M_d}$$



Newly born clumps behave like mini-starbursts


$\text{SFR} \sim 30 M_{\odot}/\text{yr}$

$M_{\star} \leq 3 \times 10^8 M_{\odot}$

$M_{\text{gas}} \leq 2.5 \times 10^9 M_{\odot}$

$\text{sSFR} > 30 \times \text{sSFR}_{\text{gal,MS}}$

$\text{SFE} > 10 \times \text{SFE}_{\text{gal,MS}}$





Clumps formation rate

Constraints on **clumps formation rate (CFR)**:


$$\text{CFR} = \frac{N_{\text{young}}}{t_V N_{\text{gal}}}$$

→ # of young clumps = 1
→ # of sample galaxies = 57
↓ Visibility window = 7 Myr




ID568


CFR ~ 2.5 clumps/galaxy/Gyr





Clumps lifetime



Genzel+ 11



Wuyts+ 13



ID568

Constraints on **clumps lifetime** (LT):

$$LT = \frac{N_{\text{cl/gal}}}{\text{CFR}} \longrightarrow \# \text{ of clumps/galaxy with } M_{\text{tot}} \geq 2.5 \times 10^9 M_{\odot}$$

LT ~ 500 Myr → clumps seem to survive stellar feedback

Summary

The birth of a star forming clump...

- Spatially resolved emission line maps of a SF galaxy at $z \sim 2$: bright off-nuclear [OIII] without a continuum counterpart
- The emission lines are powered by star formation
- It is an **extremely young star forming clump** likely formed due to violent disk instability
- It is the first direct observation of the clumps' formation phase
- **Young clumps behave like mini-starbursts** (obs. + sim.)
Old clumps have enhanced SFE (sim.)
- It supports the scenario where **clumps survive stellar feedback**

